INDEX TO VOLUME 35

New names, and the final members of new combinations, are in **bold face type.**

Acalypha gracilens, 352; virginica, 352
Acarospora **Carnegiei,** 297
Achillea, 358; Millefolium, 352
Acnida cannabina, 544; tamariscina, 544; tamariscina tuberculata, 544, tuberculata, 544
Acrolejeunea, 161, 162, 164, 165; polycarpa, 162
Acrostichum, 27
Actinopteris, 27
Additional Philippine Polyporaceae, 391
Adiantum, 27, 28; diaphanum, 272; setulosum, 272
Adopogon Dandelion, 548
Aecidium aurantiacum, 506; Botryapites, 506; germinale, 506
Agastache, 551; nepetoides, 551; scrophulariaefolia, 551
Agathis, 249–254, 256
Agoseris, 548
Agrimonia, 352, 358; Eupatoria, 552; mollis, 552
Agropyron, 533; littoreum, 200; repens, 200, 476; repens littoreum, 200; repens pilosum, 476
Agrostis alba, 191; alba aristulata, 473; alba vulgaris, 191; **antecedens,** 473–475; depressa, 192; elata, 192, 193; hyemalis, 192, 353, 474, 475; intermedia, 193; maritima, 191; perennans, 193
Aira caryophyllea, 193
Alabastra philippinensia, 63
Albertia, 250
Albugo Bliti, 362, 544; candida, 545, 546; Cladothricis, 362; **Froelichiae,** 362; Ipomoeae-panduranae, 549; Lepigoni, 543; occidentalis, 361, 548; platensis, 361, 362, 543; Portulacae, 361, 552; Tragopogonis, 361, 544, 546–549; **Trianthemae,** 361, 543
Alcicornium, 32
Algal periodicity in certain ponds and streams, 223
Alisma, 215–217, 224, 225; Plantago, 213, 217; Plantago-aquatica, 213; subcordatum, 62
Allionia incarnata, 543; nyctaginea, 543
Allium Cepa, 543
Alopecurus aristulatus, 472, 473; geniculatus, 472, 473; pratensis, 191, 473
Alsophila, 27
Amaranthus, 544; albus, 544; Bigelovii, 544; blitoides, 544; chlorostachys, 544; crispus, 544; emarginatus, 544; graecizans, 544; hybridus, 544; hybridus paniculatus, 544; Palmeri, 544; paniculatus, 544; retroflexus, 544; spinosus, 544; tristis, 544; viridis, 544
Amauroderma **asperulatum,** 407; **bataanense,** 407; **Clemensiae,** 408; Elmerianum, 408; **Ramosii,** 408
Ambrosia, 544; artemisiaefolia, 352, 544; psilostachya, 544; trifida, 352, 544
Amelanchier, 499
American botanical literature, Index to, (1907), 45; (1908), 91, 151, 219, 277; (1907), 315; (1908), 367, 417, 467, 513, 555; (1904–1907), 585
Ammophila, 56, 186; arenaria, 472
Ampelopsis quinquefolia, 544; Richii, 544; Veitchii, 544
Anabaena, 242, 247
Anacharis, 463; Alsinastrum, 461, 465; canadensis, 462, 465; Nuttallii, 461, 463, 465
Analogy between the development of the plates of crinoids and the leaves of Sassafras, 571
Anaphalis margaritacea, 352
Anaxagorea javanica, 66; **radiata** 66, 74
Anchistea virginica, 53
Andropogon, 147, 148, 353; abbreviatus, 182; furcatus, 182; littoralis, 182; virginicus, 147, 148, 182; scoparius, 148, 181; scoparius villosissimus, 181
Androsace officinalis, 552
Anemia, 27
Anemone, 552; acutiloba, 552, canadensis, 552; cylindrica, 352; dichotoma, 552; multifida, 552; nemorosa, 552; pennsylvanica, 552; quinquefolia, 552; virginiana, 552
Angiopteris, 27
Anoplolejeunea, 155, 165, 173–175; conferta 175–179; herpestica, 173, 175
Antennaria plantaginifolia, 546
Anthoxanthum odoratum, 190
Apalanthe, 463; Schweinitzii, 463, 465

Apples, The Fruit Spot of, 423
Arabis, 545; furcata, 545; glabra, 545; hirsutus, 545; Holboellii, 545; lyrata, 545; virginica, 545
Araucaria, 250–254, 256; Bidwilli, 250, 255, 257, 259, 260; **bladenensis**, 255, 257–259, 260; bohemica, 254; Cookii, 259; excelsa, 254; **Jeffreyi**, 258, 260; Toucasi, 257.
Araucarian remains from the Atlantic costal plain, 249
Araucarioxylon, 254
Araucarites, 254; ovatus, 253, 256, 260; **Zeilleri**, 252, 254, 259, 260
Archilejeunea, 155, 165, 168, 169, 171, 172, 174, 175; Auberiana, 168–171; conferta, 165, 176; Cruegeri, 168, 169; porelloides, 165, pseudo-cucullata, 165; Sellowiana, 173; **viridissima**, 169–171, 179; xanthocarpa, 165, 172
Arcyria denudata, 579; incarnata, 579
Areolaria, 295
Argemone platyceras, 363, 551
Aristida, 147, 148; dichotoma, 147, 148, 190, 353; gracilis, 190, 353; purpurascens, 148, 191
Aronia, 505, 508
Artemisia, 546; biennis, 546; canadensis, 546; ludoviciana, 546
Arthrospira, 248; Jenneri, 248
Aruncus, 541, 542; acuminatus, 542; Aruncus, 542; kamchaticus, 542; pubescens, 541; sylvestris americanus, 541
Asclepias syriaca, 352
Asparagus Sprengeri, 271, 275
Aspidium Boottii, 136; cristatum × spinulosum, 136; marginale, 18; spinulosum Boottii, 136
Asplenium, 27; ebenoides, 27, 135; Filix-foemina, 53; Filix-foemina angustum, 53; Filix-foemina Michauxii, 53; Filix-foemina rhaeticum, 53
Aster, 354, 358, 547; diffusus, 352; Novae-Angliae, 352, 546; oblongifolius, 546; sagittifolius, 547; vimineus, 352
Astilbe, 541
Astragalus, 358, 550; canadensis, 550; hastata, 548; lotiflorus, 550
Atlantic coastal plain, Some Araucarian remains from the, 249
Avena, 358
Azolla, 27

Bahamian species of Evolvulus, 89
Barbarea Barbarea, 545; vulgaris, 545
BARNHART, J. H. The published work of Lucien Marcus Underwood, 17
Basidiophora entospora, 546–548; Kellermanii, 544
Batatas edulis, 549
BERRY, E. W. Some Araucarian remains from the Atlantic coastal plain, 249
Beta vulgaris, 548
BICKNELL, E. P. The ferns and flowering plants of Nantucket, 49, 181, 471
Bicuculla, 550, 551; canadensis, 550; Cucullaria, 550
Bidens, 358; cernua, 547; chrysanthemoides, 547; comosa, 547; connata, 547; connata comosa, 547; frondosa, 343, 352, 547; laevis, 547; latifolia, 343; leucantha, 344; pilosa, 343, 344
Biographical sketch of Lucien Marcus Underwood, 1
Bjerkandera adusta, 391
Blechnum, 27
Blitum capitatum, 548
Blephariglottis Blephariglottis, 483
Boerhaavia anisophylla, 543; diffusa, 543; erecta, 543; hirsuta, 543; paniculata, 543; Sonorae, 543; spicata, 543; viscosa, 543; Xanti, 543
Boleti of the Frost herbarium, 517
Boletus affinis, 525; albus, 518; alveolatus, 518, 520; americanus, 520; ampliporus, 517; aureobrunneus, 525; auriporus, 518, 520, 521, 525; bicolor, 518, 522, 525; bovinus, 518; brevipes, 524; caespitosus, 521; canus, 525; castaneus, 518, 522; chromapes, 518; chrysenteron, 518, 524, 526; Clintonianus, 523–525; collinitus, 518; cyanescens, 519; decorus, 519; diffractus, 525; edulis, 519, 521, 522, 525; elbensis, 517; elegans, 519; eximius, 522; Farlowi, 525; felleus, 519, 521; ferrugineus, 519; firmus, 519; flavidus, 520, 521; flavo-aureus, 525; flavus, 520; Frostii, 520; glutinipes, 525; gracilis, 517, 526; graminicola, 525; graminis, 525; granulatus, 518, 520, 524; griseus, 520; innixus, 520; interruptus, 525; lenticularis, 525; limatulus, 521; luridus, 519, 521, 523, 524; luteus, 519–522; magnisporus, 521; miniato-olivaceus, 521; multipunctus, 522; Murraii, 521, 526; ornatipes, 519–522, 525; pallidus, 521; paludosus, 526; paluster, 517; Peckii, 521, 526; pictus, 521, 524, 526; piperatus, 521, 523; Ravenelii, 519, 522, 526; retipes, 519, 521, 522; robustus, 522; Roxanae, 522; rubeus, 522; rubripes, 526; Russellii, 522; salmonicolor, 522; Satanus, 523; scaber, 523, 524; serotinus, 523; sistotrema, 523; sordidus,

523, spadiceus, 523; speciosus, 523; Spraguei, 523, 524, 526; strobilaceus, 524; subchromeus, 522, 526; subreticulatus, 526; sulfureus, 522; sulfureus auricolor, 522; tenuiculus, 524; unicolor, 524; vermiculosus, 523; versipellis, 523, 524; vinaceus, 526; viridarius, 524; viscosus, 524
Botanical literature, American (1907), 45; (1908), 91, 151, 219, 277; (1907), 315; (1908), 367, 417, 467, 513, 555; (1904–1907), 585
Botrychium, 26–28, 30; obliquum, 52, 353; ternatum, 26; silaifolium, 32
Brachiolejeunea, 155, 157–159, 161, 162, 168; **bahamensis,** 383, 385, 386, 389; bicolor, 158, 159; chinantlana, 160, 386; corticalis, 158, 160, 164, 385; **densifolia,** 158, 160, 386; **insularis,** 159–161, 179, 386; latifolia, 155; sandvicensis, 157
Brachyphyllum, 250, 254
Brassica, 545; alba, 545; arvensis, 545, 546; campestris, 545; integrifolia, 545; nigra 545; oleracea, 545; sativa, 545; Sinapistrum, 545
Bremia Lactucae, 548, 549
BRITTON, N. L. Professor Underwood's relation to the work of the New York Botanical Garden, 39; Studies of West Indian plants, 337, 571; The generic name Bucida, 303; The genus Ernodea Swartz: a study of species and races, 203
Bromus, 533; hordeaceus, 200; racemosus, 200; secalinus, 200; sterilis, 200, 475; tectorum, 199, 200
BROOKS, C. The Fruit Spot of apples, 423
BROWN, H. B. Algal periodicity in certain ponds and streams, 223
Brunella vulgaris, 360
Bryonopsis laciniosa erythrocarpa, 549
Bryopteris, 157, 162
Buceras, 303
Bucida, 303; Buceras, 303, 305
Bucida, The generic name, 303
Bucida Buceras, The hypertrophied fruit of, 305
Bulbochaete, 234, 245; crenulata, 237, 245
Bursa Bursa-pastoris, 545
Bursera, 341; angustata, 342; glauca, 342; gummifera, 341; gummifera glabra, 341; gummifera pubescens, 341; inaguensis, 342; simplicifolia, 342

Cactus divaricatus, 564; repandus, 564
Caeoma, 506; Botryapites, 506; germinale, 506
Cakile americana, 545; edentula, 545
Caladium Colocasia, 544
Calamagrostis canadensis, 472; cinnoides, 472
Calandrinia Menziesii, 552
Callipteris, 27
Callitriche, 224
Calonyction aculeatum, 549
Calophyllum Inophyllum, 400
Caloplaca **amabilis,** 298; elegans **brachyloba,** 298
Calothrix parietina, 243, 248
Calvatia, 295; sculptum, 295
Calystegia Sepium, 549
Camelina microcarpa, 545; sativa, 545
Campanula, 358
Camptosorus, 27
Cantharellus, 27
Capnoides, 551; aureum, 550, 551; sempervirens, 550, 551
Capsella Bursa-pastoris, 545
Cardamine bulbosa, 545; hirsuta, 352, 545; laciniata, 545; ludoviciana, 545; rhomboidea, 545
Carduus arvensis, 547; lanceolatus, 547; muticus, 547; spinosissimus, 547; undulatus, 547
Carex, 262, 266, 267, 358, 471, 487, 490, 494; **abdita,** 492; adusta, 498; albicans, 490; albolutescens, 497; alpestris, 489; annectens, 492; aquatilis, 487; brunnescens, 496; bullata, 484; bullata Greenii, 484; canescens, 496; canescens alpicola, 496; canescens disjuncta, 495; canescens subloliacea, 496; **cephalantha,** 489, 493–495; cephalantha angustata, 494; **chihuahuensis,** 265; communis, 490; comosa, 485; costellata, 488, 489; costata, 488; cristata, 264; Davisii, 490; deflexa, 490; delicatula, 495; disjuncta, 495; echinata cephalantha, 493; Emmonsii, 489–491; festiva, 262; festucacea, 353; feta, 269; filiformis, 485, 486; **gigas,** 266–268; Goodenovii, 486–489; gynandra, 488; hirta, 486; hormathodes, 496, 497; hormathodes invisa, 496, 497; **incomperta,** 494; interior capillacea, 495; intumescens, 483; lagopina, 262; lagopodioides moniliformis, 264; lanuginosa, 485, 486; leporina, 261, 262; leporina americana, 261; leptalea, 492; Liddoni, 263, 264; lupulina, 484; lupulina pedunculata, 484; lurida, 353, 484; lurida flaccida, 484; marcida, 266; Muhlenbergii, 353, 493; muricata, 493; pallescens, 353, 489; pennsylvanica, 489; petasata, 261–263; phaeocephala, 261–263; physaema, 484; Preslii, 261, 262;

projecta, 264; pseudo-cyperus, 485; pseudoscirpoidea, 266–268; reducta, 264; rosea, 493; **scabriuscula,** 267, 286; **scirpiformis,** 267, 268, 270; scirpoidea, 266, 267, 269, 270; scirpoidea gigas, 268; scirpoidea stenochlaena, 269; scirpoides capillacea, 495; seorsa, 495; silicea, 497; stellulata cephalantha, 493; stellulata excelsior, 493; **stenochlaena,** 266, 267, 269; sterilis, 494, 495; stipata, 492; straminea, 496, 497; straminea alata, 498; straminea echinodes, 496, 497; stricta, 486; subloliacea, 496; tenuis, 489; **tonsa,** 492; tribuloides, 265; tribuloides moniliformis, 264; umbellata, 491, 492; umbellata brevirostris, 491; umbellata tonsa, 491, 492; utriculata, 484; utriculata minor, 484; varia, 490, 491; vestita, 485; virescens, 488, 489; virescens minima, 488; virescens Swanii, 488; vulgaris strictiformis, 487; vulpinoidea, 492; Walteriana, 485; xanthocarpa annectens, 492
Carex, Notes on, 261
Carica Papaya, 141, 145
Carica Papaya, Correlation of flower- and fruit-structure in, 141
Cassia, 397; marylandica, 352
Cassipourea, 339; alba, 339; elliptica, 339, 340; guianensis, 339; **subcordata,** 340; **subsessilis,** 340
Castalia ampla, 214; odorata, 217, 283, 284, 286, 287–289
Caudalejeunea, 162
Celastrus scandens, 352
Celtis occidentalis, 553
Cenchrus tribuloides, 190
Centaurea, 358, 547
Cephalocereus, 566
Cephalozia, 25
Cerastium, 543; cerastioides, 543; longipedunculatum, 543; nutans, 543; trigynum, 543; viscosum, 543; vulgatum, 543
Ceratites Botryapites, 506
Ceratopteris, 27; triangularis, 31
Cereus aggregatus, 84; coccineus, 84; conoideus, 86; cubensis, 562; divaricatus, 565; divergens, 565; erectus, 566; eriophorus, 562; gracilis, 563; monoclonos, 565; pellucidus, 562; phoeniceus, 84; phoeniceus conoideus, 86; polyacanthus, 85; repandus, 563, 564, 566; Roemeri, 86; subrepandus, 563; undatus, 564
Chaetochloa, 353, 552; glauca, 190, 552; italica, 190, 552; versicolor, 190; viridis, 190, 552
Chaetophora, 237–239, 244, 246; endiviaefolia, 239, 246; pisiformis, 229, 235, 238, 246
Chamaecrista, 352, 358
Chamaecyparis, 501; nootkatensis, 501; thyoides, 510, 511
Chamaeraphis, 552
Chamaesyce, 550; cordifolia, 550; glyptosperma, 550; hirtula, 550; humistrata, 550; serpens, 550; serpyllifolia, 550; stictospora, 550
Cheilanthes, 27
Cheilolejeunea, 378
Cheiranthus asper, 545; Cheiri, 545; pacificum, 545
Chenopodium, 548; album, 548; album viride, 548; hybridum, 548; leptophyllum, 548; rubrum, 548
Chloris elegans, 552
Chrysomyxa, 499
Cibotium, 27
Cichorium Intybus, 548
Cienkowskia reticulata, 577
Cinnamomum Heerii, 575
Cirsium, 358; arvense, 547
Citrullus vulgaris, 549
Cladium mariscoides, 483
Cladophora, 233, 234, 240, 246; glomerata, 229, 240, 244, 246
Cladothrox lanuginosa, 362, 544
Claviceps, 529, 533; purpurea, 533
Claytonia perfoliata, 552; virginica, 552
Clethra **Williamsii,** 73, 75
Closterium, 109, 110, 133, 232, 245; abruptum, 124; acerosum, 121, 227, 232–234, 245; acerosum elongatum, 122, 134; aciculare, 128, 134; acuminatum, 117; acutum, 128, 134; angustatum, 113; angustatum clavatum, 113, 133, 134; angustatum decussatum, 113; angustatum reticulatum, 113; Archerianum, 111, 134; arcuatum, 118; areolatum, 130; attenuatum, 127; Braunii, 130; Calamus, 131; calosporum, 118, 119; Cornu major, 124; costatum, 114, 134; costatum Westii 114, 134; Cucumis, 120, 245; Cynthia, 110, 133; decorum, 131, 134; decussatum, 133; Delpontei, 131; Dianae, 117, 118, 134, 245; Dianae arcuatum, 118; didymotocum, 111, 112, 134; didymotocum **Johnsonii,** 112; eboracense, 120; Ehrenbergii, 121, 134, 233, 245; Ensis, 111; gracile, 126; gracile elongatum, 126; gracile tenue, 126; incurvum, 119, 134; intermedium, 115; intervalicola, 131, 134; Jenneri, 119, 134; Johnsonii, 112; juncidum, 116, 134; juncidum brevius, 117, 134; juncidum elongatum, 117; Kützingii, 132, 134; lanceolatum, 122; Leibleinii, 120, 134;

Leibleinii curtum, 121; limneticum tenue, 126; lineatum, 129, 134; lineatum costatum, 129; littorale, 124, 134; Lunula, 122, 134; Lunula intermedium, 123; Lunula maximum, 123, 134; Lunula minor, 123; Lunula minus, 123; macilentum, 112, 134; maculatum, 130; Malmei, 111, 134; moniliferum, 121, 134; mourense, 131; nasutum, 133; **Novae-Angliae,** 131, 134; parvulum, 118, 134; praelongum, 125; Pritchardianum, 127, 134; pronum, 127, Pseudodianae, 118, 134; Ralfsii, 129; Ralfsii hybridum, 130, 134; Ralfsii immane, 130, 134; regulare, 114, 134; robustum, 121; rostratum, 132, 134; rostratum brevirostratum, 132; setaceum, 133, 134; strigosum, 125, 245; striolatum, 115, 134; striolatum elongatum, 116; striolatum erectum, 115, 134; striolatum intermedium, 115; subangustatum, 133; subangustatum clavatum, 133; subdirectum, 116; subtile, 126; subulatum, 128, 134; Toxon, 125, 134; tumidum, 124; turgidum, 127; turgidum intermedium, 127; Ulna, 116, 134; Venus, 119, 134

Closterium, The New England species of, 109

Cnicus, 547; arvensis, 547; horridulus, 547; lanceolatus, 547; muticus, 547

Coccinia indica, 549

Cochlearia Armoracia, 545; oleracea, 545

Coelastrum, 241, 247; microsporum, 247

Coleochaete, 245; scutata, 238, 245

Coleosporium Campanulae, 25

Colocasia Antiquorum, 544

Color variation in some of the fungi, 307

Coltricia **benguetensis,** 391

Columbea, 250, 253

Conocephalum, 325

Convolvulaceae, Studies in the North American, 97

Convolvulus, 549; altissimus, 99; Batatas, 549; bractiflorus, 100; eriospermus, 105; filiformis, 104; incanus, 549; macrorrhiza, 549; obvallatus, 100; racemosus, 99; repandus, 104; Sepium, 352, 549

Cook, M. T. The development of the embryo-sac and embryo of Potamogeton lucens, 209; The hypertrophied fruit of Bucida Buceras, 305

Cordyceps, 529, 532; militaris, 532; pistillariaeformis, 532

Coreopsis leucantha, 344

Coriolopsis aneba, 407; badia, 392; **bataanensis,** 393; caperata, 392; **Copelandi,** 392; dermatodes, 393; **melleoflava,** 393; occidentalis, 394; semilaccata, 394; **subcrocata,** 394

Coriolus atypus, 394; **Clemensiae,** 394; **Currani,** 395; elongatus, 395; maximus, 395; murinus, 395; nigromarginatus, 396; **perpusillus,** 396; prolificans, 396; **rubritinctus,** 396; **subvernicipes,** 397; versicolor, 403, 531

Cornucopiae hyemalis, 192

Cornus candidissima, 352

Coronopus, 545

Correlation of flower-and fruit-structure in Carica Papaya, 141

Corydalis, 551; aurea occidentalis, 551; glauca, 551

Cosmarium, 109, 234, 245; Botrytis, 245; laeve, 245; Phaseolus, 245; pyramidatum, 245

Crataegus, 350, 352, 354–357, 583; coloradensis, 581–583; macracantha occidentalis, 581, 582; ocidentalis, 581–583; spathulata, 503, 504

Crataegus, Two imperfectly known species of, 577

Crepis, 358

Crinoids, An analogy, 571

Crocus, 358

Crossotolejeunea, 373

Cryptogramma, 27

Cucumis angulatus, 549; Anguria, 549; dipsaceus, 549; erinaceus, 549; Melo, 549; odoratissimus, 549; sativus, 549

Cucurbita esculenta, 550; maxima, 550; ovifera, 550; Pepo, 550; verrucosa, 550

Cunninghamia, 253

Cunninghamites, 253; elegans, 253; squamosus, 253

Curtis, C. C. A biographical sketch of Lucien Marcus Underwood, 1

Cushman, J. A. The New England species of Closterium, 109

Cyathea, 27, 31

Cyathocalyx **acuminatus,** 66, 74; biovulatus, 67

Cyathula lappulacea, 544

Cyclomyces fuscus, 414

Cycloporellus **barbatus,** 397; cichoriaceus, 397; microcyclus, 397

Cylindrosporium, 452; **Pomi,** 446, 453; Ranunculi, 452

Cynoglossum officinale, 545

Cyperus anceps, 568; bromoides, 568; dentatus, 476; dentatus ctenostachys, 476; diandrus, 476; digitatus, 568; erythrorhizos, 478; esculentus, 477.

568; esculentus angustispicatus, 477; esculentus leptostachyus, 477; filiculmis, 353, 478; filiculmis macilentus, 478; granularis, 568; Grayi, 478; humilis, 568; **macilentus,** 478; Nuttallii, 476; oxylepis, 568; rivularis, 476; speciosus, 477; strigosus, 478; strigosus compositus, 478; strigosus robustior, 478; tetragonus, 568
Cyrtolejeunea, 165
Cyrtomium, 27
Cystopteris, 27

Dactylis glomerata, 196
Daedalea amanitoides, 405, 414; Hobsoni, 414; **isabellina,** 414; **subconfragosa,** 415
Dammara, 249, 250, 254, 256
Dammarites, 249
Danaea, 30; alata, 382
Danthonia Alleni, 193; compressa, 475; Faxoni, 193; sericea, 193; spicata, 193
Daphne **luzonica,** 72, 75
Davallia, 27
Dennstaedtia, 27; punctilobula, 54
Dentaria, 546; diphylla, 545; heterophylla, 545; laciniata, 545; maxima, 545
Deparia, 27
Deschampsia flexuosa, 193
Development of the embryo-sac of Nymphaea advena, 283
Development of the embryo-sac and embryo of Potamogeton lucens, 209
Development of the sexual organs and sporogonium of Marchantia polymorpha, 321
Dianthus, 358
Dibrachiella, 168, 169
Dicksonia, 27
Dicranolejeunea, 178
Dictydiaethalium plumbeum, 578
Dictydium cancellatum, 578
Dictyogramma, 27
Diderma crustaceum, 577
Didymochlaena, 27
Didymium squamulosum, 578
Dicentra, 551
Diplachne **maritima,** 195; procumbens, 195
Diplazium, 27
Dipsacus sylvestris, 550
Dipterocarpus, 409
Distichlis spicata, 196
Docidium, 245; Trabecula, 234, 245
DODDS, G. S., & RAMALEY, F. Two imperfectly known species of Crataegus, 577
Dorstenia **jamaicensis,** 567
Doryopteris, 27
Dothidea pomigena, 425
DOWELL, P. New ferns described as hybrids in the genus Dryopteris, 135
Draba caroliniana, 546
Draparnaldia, 237, 238, 246; plumosa, 229, 239, 244, 246
Drosera filiformis, 55, 483
Drymoglossum, 27
Drynaria, 27
Dryopteris, 27, 28, 135; Boottii, 54, 136; Clintoniana, 137; **Clintoniana × Goldiana,** 137; **Clintoniana × intermedia,** 136; cristata, 54, 135–137; **cristata × intermedia,** 136, 137; cristata × marginalis, 135; cristata × spinulosa, 136; Filix-mas × spinulosa, 135; Goldiana, 139; Goldiana celsa, 137, 138; **Goldiana × intermedia,** 138; **Goldiana × marginalis,** 139; intermedia, 54, 137, 139; marginalis, 135, 139; marginalis × spinulosa, 135; noveboracensis, 53; pittsfordensis, 135; simulata, 53, 54; spinulosa, 54, 136; Thelypteris, 53
Dryopteris, New ferns described as hybrids in the genus, 135
Dulichium arundinaceum, 479
DURAND, E. J. The development of the sexual organs and sporogonium of Marchantia polymorpha, 321

Earliella corrugata, 398
Echinella acuta, 128
Echinocerei of New Mexico, 77
Echinocereus coccineus, 77–82, 84; conoideus, 78–81, 84–86; **neo-mexicanus,** 84, 86, 87; phoeniceus conoideus, 86; polyacanthus, 78–81, 83–85
Echinochloa Crus-galli, 183; Crus-galli mutica, 183; Walteri, 183
Echinocystis lobata, 550
Echinospermum Redowskii cupulatum, 545; Redowskii occidentale, 545
Elaeocarpus tuberculatus, 72; **venosus,** 71, 75
Eleocharis, 224, 225, 236; acicularis, 480; capillacea, 569; glaucescens, 479, 480; obtusa, 353, 479; palustris, 479, 480; rostellata, 480; tenuis, 480; tricostata, 480
Elfvingia Elmeri, 409; tornata, 409
Elodea, 457; canadensis, 457–461, 463, 465; latifolia, 463, 465; Planchonii, 462, 465; Schweinitzii, 463, 465
Elymus **halophilus,** 201; hirsutiglumis, 201; virginicus, 201
Embryo-sac of Nymphaea advena, The development of the, 283
Embryo-sac and embryo of Potamo-

geton lucens, The development of the, 209
Encrinus liliiformis, 571, 572
Epicostorus, 535
Equisetum, 24; arvense, 54, 353; fluviatile, 54; variegatum, 19
Eragrostis major, 195; pectinacea, 195, 353; Purshii, 195, 353; spectabilis, 196
Erechtites hieracifolia, 547
Erigeron, 547; annuus, 547; canadensis, 547; philadelphicus, 547; ramosus, 352, 547
Eriogynia, 541
Eriophorum gracile, 482; tenellum, 482; virginicum, 482; virginicum album, 482; viride-carinatum, 482
Eriophyes, 305, 306
Ernodea, 203; angusta, 203, 204, 206, 207; Cokeri, 203, 204, 207, 208; littoralis, 203, 204, 206–208; **Millspaughii,** 204, 207; **Nashii,** 204, 208; **Taylori,** 204, 208
Ernodea Swartz, The genus: a study of species and races, 203
Erysimum, 358
Eucaloplaca, 298
Euglena viridis, 227
Euleptogium, 299
Euosmolejeunea, 378
Eupatorium ageratoides, 547; purpureum, 547
Euphorbia, 358, 363, 550; glyptosperma, 363; hirsuta, 550; hypericifolia, 550; maculata, 363, 550; Preslii, 352, 550
Euphrasia, 358
Eutacta, 250, 253, 259
Euthamia caroliniana, 352; graminifolia, 352
EVANS, A. W. Hepaticae of Puerto-Rico, 155; New West Indian Lejeuneae, 371
Evolvulus Arbuscula, 89; **bahamensis,** 89; **Bracei,** 90; incanus, 90; purpureo-coeruleus, 89; squamosus, 89; Wrightii, 90
Evolvulus, Two Bahamian species of, 89
Exogonium, 97; arenarium, 98, 104; argentifolium, 97, 98, 102; bracteatum, 98, 100; bracteatum **pubescens,** 101; **Conzattii,** 98, 102; **cubense,** 98, 105; **Eggersii,** 98, 104; eriospermum, 98, 105; filiforme, 98, 104; **fuchsiodes** 98, 101; **jalapoides,** 98, 101; **leuconeurum,** 98, 106; **luteum,** 98, 103; **microdactylum,** 98, 101, 102; microdactylum **integrifolium,** 103; Olivae, 100; pedatum, 98, 106; racemosum, 98, 99; repandum, 98, 104; Rudolphii, 98, 99; spicatum, 100; **velutifolium,** 98, 100; **viridiflorum,** 98, 106; **Wrightii,** 98, 99
Exogonium, The genus, 97

Favolus philippinensis, 401; **resinosus,** 398; **subrigidus,** 398; tenuis, 399; transiens, 397; Wightii, 399
Ferns described as hybrids in the genus Dryopteris, New, 135
Ferns and flowering plants of Nantucket, 49, 181, 471
Festuca, 358; capillata, 198; duriuscula, 199, 475; elatior, 199; Myuros, 198; octoflora, 197; ovina, 198, 199, 475; ovina duriuscula, 199; pratensis, 199; rubra, 198, 199
Fimbriaria, 324
Fimbristylis autumnalis, 353, 569
Floerkea proserpinacoides, 551
Flower- and fruit-structure in Carica Papaya, Correlation of, 141
Flowering plants of Nantucket, 49, 181, 471
Fomes Auberianus, 409; ligneus, 409; luzonensis, 409; pyrrhocreas, 411; **subresinosus,** 410; substygius, 412; **subungulatus,** 410; ungulatus, 409; Warburgianus, 407
Fossombronia, 24
Fragaria americana, 352; virginiana, 352
Franseria discolor, 544; tenuifolia, 544
Freycinetia Cumingiana, 64; luzonensis, 64
Froelichia campestris, 362, 544; floridana, 362, 544; gracilis, 362, 544
Frost herbarium, The Boleti of the, 517
Fruit of Bucida Buceras, The hypertrophied, 305
Fruit Spot of apples, 423
Fruit-structure in Carica Papaya, Correlation of flower- and, 141
Frullania, 157, 385, 386
Frullanoides densifolia, 158, 159
Frustularia subulata, 128
Fuligo ovata, 578
Fumaria, 358
Funalia funalis, 399; leonina, 399; philippinensis, 399; villosa, 399
Fungi, Color variation in some of the, 307
Fusarium, 529
Fusicladium dendriticum, 424

Gaertneria acanthocarpa, 544; discolor, 544

Galium, 358, 553; Aparine, 553; boreale, 553; lanceolatum, 553; triflorum, 553
Ganoderma amboinense, 410; **balabacense,** 410; **Currani,** 411; subtornatum, 409
Gasparrinia, 298
Gaura parviflora, 551
Generic name Bucida, The, 303
Genus Dryopteris, New ferns described as hybrids in the, 135
Genus Ernodea Swartz: a study of species and races, The, 203
Genus Exogonium, The, 97
Genus Gymnosporangium, Studies in the, 499
Geranium, 551; carolinianum, 551; dissectum, 551; maculatum, 551; pusillum, 551; Richardsonii, 551; Robertianum, 551
Geum album, 552; canadense, 552; macrophyllum, 553; rivale, 553
Gibberella, 528, 529; pulicaris, 529
Gilia, 552
Gleichenia, 28
Gloecystis, 247; gigas, 247
Gloeophyllum edule, 415; **nigrozonatum,** 415; striatum, 416
Glyceria fluitans, 196; septentrionalis, 196
Gnaphalium obtusifolium, 352; purpureum, 547; spathulatum, 547
Gomphostrobus, 250
Goniopteris, 28
Gonium, 241, 246; pectorale, 246
Gonolobus, 544; macrophyllus, 544; suberosus, 544
Grasses from the West Indies, Two new, 301
Grossularia divaricata, 551; oxyacanthoides, 551; rotundifolia, 551
Gymnogramma, 28
Gymnogramme, 30
Gymnosporangium, 25, 499–502, 504, 505; aurantiacum, 509; Betheli, 510; bermudianum, 502, 509; biseptatum, 506; Botryapites, 506, 510; clavariaeforme, 504, 510; clavipes, 506; **Davisii,** 507, 511; durum, 510; Ellisii, 505, 510, 511; **exiguum,** 508, 511; flaviforme, 504; flaviformis, 504; **floriforme,** 503, 504, 510; floriformis, 504; germinale, 506, 510; globosum, 510; inconspicuum, 508, 511; juniperinum, 505, 510; Juniperi-virginianae, 502; **Libocedri,** 509, 511; macropus, 502; Nelsoni, 508, 511; Nidus-avis, 510, 511; speciosum, 505, 510; tremelloides, 505
Gymnosporangium, Studies in the genus, 499
Hapalopilus gilvus, 399; licnoides, 399; malaiensis, 400; **Ramosii,** 400; subrubidus, 400
Harpalejeunea, 375, 377–379; **reflexula,** 375, 376, 389; subacuta, 377, 379, 380; uncinata, 376, 377
HARPER, R. M. Some native weeds and their probable origin, 347
Harrisia, 561; **Brookii,** 562, 564; **eriophorus,** 561, 562; **Fernowi,** 561, 562; **gracilis,** 561, 564; **Nashii,** 562, 564; **portoricensis,** 561, 563; **Taylori,** 562, 565; **undata,** 562, 564
HARSHBERGER, J. W. The water-storing tubers of plants, 271
Hartmannia speciosa, 551
Hedeoma hispida, 551
Helianthus, 547; annuus, 547; divaricatus, 547; doronicoides, 547; grosseserratus, 547; hirsutus, 547; occidentalis, 547; Maximiliani, 547; scaberrimus, 547; strumosus, 547; trachelifolius, 547; tuberosus, 547
Hemionitis, 28
Hemitelia, 28
Hemitrichia clavata, 579; vesparia, 579
Heppia **placodizans,** 299; **desertícola,** 300; Hassei, 300; leptopholidis, 300
Hepatica acuta, 552; acutiloba, 552; Hepatica, 552; triloba, 552
Hepaticae of Puerto Rico, 155
Hernandia, 337, 338; cubensis, 338; guianensis, 338; **jamaicensis,** 338; sonora, 338; sonora guadeloupensis, 338
Herposteiron, 238, 245
Hesperis matronalis, 546
Hexagona cucullata, 400; **luzonensis,** 401; **pertenuis,** 401; philippinensis, 401
Hieracium, 358
Holcus lanatus, 193
Homalocenchrus oryzoides, 190
Hordeum jubatum, 476; vulgare, 201
HOUSE, H. D. Studies in the North American Convolvulaceae, 97; Two Bahamian species of Evolvulus, 89
Houstonia coerulea, 352; minor, 553; patens, 553
HOWE, M. A. Lucien Marcus Underwood: a memorial tribute, 13
Humata, 28
HUMPHREYS, E. W. An analogy, between the development of the plates of crinoids and the leaves of Sassafras, 577
Hyalotheca, 234, 245
Hybrids in the genus Dryopteris, New ferns described as, 135

Hydrophyllum macrophyllum, 551; virginicum, 551
Hymenophyllum, 28
Hyoscyamus, 364
Hypericum maculatum, 352; mutilum, 352
Hypertrophied fruit of Bucida Buceras, 305
Hypocrea, 311, 528, 532; apiculata, 311, 313; chlorospora, 310, 311, 313; citrina, 313, 532; gelatinosa, 310, 311, 313; patella, 532; Richardsoni, 532; rufa, 532; tuberculariformis, 530; viridis, 311
Hypocreales, Some North Dakota, 527
Hypolepis, 28
Hypomyces, 528, 531; aurantius, 531; lactifluorum, 312, 313, 531; ochraceus, 531; polyporinus, 531; purpureus, 312, 313, 531; rosellus, 531

Ibidium Beckii, 52
Icotorus, 535
Ilysanthes gratioloides, 352
Impatiens, 545; aurea, 545; biflora, 545; fulva, 545; pallida, 545
Index to American botanical literature (1907), 45; (1908), 91, 151, 219, 277; (1907), 315; (1908), 367, 417, 467, 513, 555; (1904–1907), 585
Inonotus **Clemensiae,** 401; fruticum, 402
Ipomoea, 97; alterniflora, 105; altissima, 99; arenaria, 104; argentifolia, 102; Batatas, 549; bracteata, 97–100, bracteata pubescens, 101; Carolina, 549; cincta, 100; commutata, 549; Conzattii, 102; Desrousseauxii, 105; eriosperma, 105; filiformis, 104; fuchsioides, 101; fuchsioides glabra, 101; fuchsioides parviflora, 101; hederacea, 549; incarnata, 549; jalapoides, 101; lacunosa, 549; leptophylla, 549; leuconeura, 106; mexicana, 549; microdactyla, 102; nematoloba, 107; obtusata, 105; pandurata, 549; Pes-caprae, 549; praecox, 102; Purga, 97; purpurea, 549; racemosa, 99; repanda, 103, 104; rubrocincta, 106; Rudolphii, 99; spicata, 100; Steudeli, 104; tamnifolia, 549; trichocarpa, 549; triloba, 549; viridiflora, 106
Isactis caespitosa, 243, 248
Isaria farinosa, 532
Isidorea, 203
Isnardia palustris, 352
Isoetes, 20, 55; echinospora Braunii, 55
Iva ambrosiaefolia, 544; ciliata, 547; xanthifolia, 544
Ixophorus, 552

Jackson, H. S. Sorosporium Ellisii Winter, a composite species, 147
Jacquemontia tamnifolia, 549
Juncus, 358; bufonius, 353; effusus, 353; marginatus, 353; tenuis, 353
Jungermannia bicolor, 158, 159; conferta, 175; polycarpa, 162; xanthocarpa, 175
Juniperus, 501; communis, 353, 510; communis alpina, 507; communis canadensis, 353; nana, 507; sabinoides, 508; sibirica, 507, 510, 511; virginiana, 58, 503, 508

Kelseya, 540, 541
Kern, F. D. Studies in the genus Gymnosporangium, 499
Kneiffia pumila, 352
Knoxia, 204
Koellia virginiana, 352
Koniga maritima, 546
Krigia Dandelion, 548; virginica, 352
Kyllinga intermedia, 568; peruviana, 568

Literature, American Botanical (1907), 45; (1908), 91, 151, 219, 277; (1907), 315; (1908), 367, 417, 467, 513, 555; (1904–1907), 585
Lobelia inflata, 352
Lolium italicum, 200, 476; perenne, 200, 475
Lomaria, 28
Lophanthus, 551
Lopholejeunea, 162
Lorinseria areolata, 52
Loxoscaphe, 28
Lunulina monilifera, 121
Luetkea, 541
Lutkea, 541
Lycogala epidendrum, 579
Lycoperdon, 293, 295; caelatum, 295; insculptum, 291; sculptum, 293, 294
Lycoperdon sculptum Harkness, Notes on, 291
Lycopersicon Lycopersicon, 553
Lycopodium, 28, 32; adpressum, 55; alopecuroides, 55, 483; complanatum, 55; inundatum Bigelovii, 55; obscurum, 55; tristachyum, 29, 55
Lyngbya, 247; Juliana, 247; ochracea, 247
Lenzites **Clemensiae,** 416; **submurina,** 416
Lepidium campestre, 546; densiflorum, 546; incisum, 546; intermedium, 546; sativum, 546; virginicum, 352, 546
Leptilon canadense, 352, 547
Leptogium **arizonicum,** 299

Lespedeza, 358, 352
Leucolejeunea, 155, 165, 171, 174, 178; clypeata, 171, 177; unciloba, 172, 173; xanthocarpa, 172, 173, 179; conchifolia, 173
Leucoporus hirto-lineatus, 401
Leucostegia, 28
Libocedrus, 501, 509; decurrens, 511
Lichens, New North American, 297
Limnocharis, 215; emarginata, 214, 217
Linaria canadensis, 352, 553
Lindsaea, 33
Linum sulcatum, 551
Liriodendron, 575
Lactaria, 312
Lactuca altissima, 548; canadensis, 352, 548; hirsuta, 548; integrifolia, 548; leucophaea, 548; ludoviciana, 548; pulchella, 548; sagittifolia, 548; sativa, 548; spicata, 548, 549
Laetiporus speciosus, 402
Lagenaria vulgaris, 550
Lamium amplexicaule, 551
Laportea canadensis, 553
Lappula cupulata, 545; floribunda, 545; occidentalis, 545; texana, 545
Larix decidua, 57
Lastraea, 28
Lechea maritima, 52
Legnotis, 339; elliptica, 339; Cassipourea, 339
Leiolejeunea, 377, 378; **grandiflora**, 378, 379, 389
Lejeunea, 21, 157, 162, 377, 383, 385; asperrima, 375; atroviridis, 165; bicolor, 159; conferta, 174, 175; conferta Miquelii, 177; conferta Liebmaniana, 177; cyclostipa, 169; domingensis, 162, 164; florentissima, 169, 171; herpestica, 175; holostipa, 165; involutiloba, 175; linguaefolia, 164; Miquelii, 175; polycarpa, 162; proteoides, 175; stricta, 377; subaurita, 175; unciloba, 169, 173; viridissima, 169; xanthocarpa, 172
Lejeuneae, New West Indian, 371

MACKENZIE, K. K. Notes on Carex, 261
Madia sativa, 547
Malache, 344; scabra, 344, 345; **troyana**, 345
Malus Malus, 453
Mammillaria aggregata, 84
Marattia, 28
Marchantia, 322–325, 327, 328; polymorpha, 322, 332
Marchantia polymorpha, The development of the sexual organs and sporogonium of, 321
Marchesinia, 155, 158
Marsilia, 19
Mastigolejeunea, 156, 161, 168
Matricaria matricariodes, 547
Matteuccia, 28
Medicago sativa, 550
Megapterium missouriense, 551
Melothria scabra, 550
Memorial tribute: Lucien Marcus Underwood, 13
Meniscium, 28
Mentha, 360; canadensis, 352
Merismopedia, 248; convoluta, 248
Mesocarpus parvulus, 231; radicans, 231; scalaris, 232
Micrampelis echinata, 550; lobata, 550
Microlepia, 28
Microporellus dealbatus, 402; subdealbatus, 402
Microzamia dubia, 253, 260
Mitrephora ferruginea, 67; Lanotan, 67; **Merrillii**, 67; **Williamsii**, 68, 74
Mohria, 28
Momordica balsamina, 550; Charantia, 550
Monoon, 68
Monotropella, 168
Montia perfoliata, 552
Mougeotia, 231, 244; genuflexa, 232, 244; parvula, 232, 244; scalaris, 232, 244
Moulds, North Dakota slime-, 577
Muhlenbergia mexicana, 191, 353; diffusa, 191
Mukia scabrella, 550
Mulgedium leucophaeum, 549
MURRILL, W. A. Additional Philippine Polyporaceae, 391; The Boleti of the Frost herbarium, 517
Myosotis verna, 545
Myxonema, 238, 239, 246; nanum, 239, 243, 246; tenue, 239, 246

Nabalus albus, 549; altissimus, 548, 549
Nageiopsis, 256, 257; montanensis, 256, 257; ovata, 256, 257; zamioides, 257
Naias, 60, 215, 217; flexilis, 60, 214; guadalupensis, 60, 61; microdon, 60
NASH, G. V. Two new grasses from the West Indies, 301
Nantucket, The ferns and flowering plants of, 49, 181, 471
Nasturtium, 546
Native weeds and their probable origin, 347
Nectria, 528, 529; cinnabarina, 308, 530; coccinea, 529; episphaeria, 529; Meliae, 310, 313; nigrescens, 309, 313; offuscata, 309, 313; Peziza, 529; purpurea, 308–310, 313, 530, 531;

Russellii, 309, 313; sanguinea, 529; tubercullariformis, 528, 529; verrucosa, 530
Neillia opulifolia multiflora, 536
Nelumbo, 283, 284, 289
Nephrocytium, 247; Agardhianum, 247
Nephrodium, 28; esculentum, 272
Nephrolepis, 28, 272–274; acuminata, 271; cordifolia, 271–273, 275; cordifolia etuberosa, 273; cordifolia tuberosa, 273; davallioides, 271, 273; exaltata, 271, 272; hirsutula, 272, 273; pectinata, 272; philippinensis, 271–273; Pluma, 272, 273; Pluma Bausei, 272; tuberosa, 271, 272, 275; undulata, 272
Neslia paniculata, 546
New England species of Closterium, 109
New ferns described as hybrids in the genus Dryopteris, 135
New grasses from the West Indies, 301
New Mexico, Some Echinocerei of, 77
New North American lichens, 297
New West Indian Lejeuneae, 371
New York Botanical Garden, Professor Underwood's relation to the work of the, 39
Nicotiana glauca, 364, 553; longiflora, 364
Nigrofomes melanoporus, 411
Nigroporus durus, 402; vinosus, 402
Niphobolus, 29
North American Convolvulaceae, 97
North American lichens, New, 297
North American Peronosporales, 361, 543
North Dakota Hypocreales, 527
North Dakota slime-moulds, 577
Nostoc, 242, 247; commune, 242, 247; verrucosum, 242, 243, 247
Notes on Carex, 261
Notes on Lycoperdon sculptum Harkness, 291
Notes on Philotria Raf., 457
Notes on Rosaceae, 535
Notholaena, 28
Nymphaea, 213, 284, 289; advena, 217, 283, 284, 289
Nymphaea advena, The development of the embryo-sac of, 283
Odontolejeunea, 381, 382; **longispica,** 380–383, 389; lunulata, 381–383; Sieberiana, 381–383
Oedogonium, 234, 237, 245; crassiusculum, 235, 237, 238, 245; cryptoporum, 227, 236, 245
Oenothera, 355, 551; biennis, 352, 551; laciniata, 551; missouriensis, 551; sinuata, 551; speciosa, 551
Oligonema flavidum, 580
Onagra biennis, 551
Onoclea, 28; sensibilis, 54; sensibilis obtusilobata, 18
Onychium, 28
Ophiocytium, 247; bicuspidatum, 247
Ophioglossum, 28; arenarium, 52; vulgatum, 51, 52
Panicum, 186–188, 353, 354, 358; agrostoides, 184; anceps, 184; atlanticum, 189; Bicknellii, 186; boreale, 189; capillare, 184; clandestinum, 189; Clutei, 189; columbianum, 187; depauperatum, 184–186; dichotomum, 186; filiculme, 187, 188; **Grisebachii,** 301; lineariifolium, 185, 186; martinicense, 301; mattamusketense, 189; meridionale, 187, 188; miliaceum, 184; oricola, 187; **Owenae,** 185; proliferum, 184; Scribnerianum, 189; sphaerocarpon, 189; tennesseense, 188; unciphyllum, 188; unciphyllum thinium, 188; virgatum, 184
Papaver, 364
Parietaria pennsylvanica, 553
Parthenium integrifolium, 547; repens, 548
Parthenocissus quinquefolia, 554; tricuspidata, 554
Paspalum Muhlenbergii, 183; prostratum, 182; psammophilum, 182; pubescens, 183; setaceum, 182
Passiflora ciliata, 343; foetida, 343
Pavonia, 345; racemosa, 344; racemosa troyana, 345; spicata, 344; spinifex, 345
Pediastrum, 241, 246; Boryanum 246
Pellaea, 28
Peltolejeunea, 178
Penium, 245; interruptum, 234, 245
Peperomia, 567
Opulaster, 535, 536, 537; alabamensis, 536, 537; alternans, 538; australis, 536, 537; bracteatus, 536; bullatus, 535; capitatus, 536, 537; cordatus, 536, 537; Hapemanii, 536–538; intermedius, 536, 537; malvaceus, 536–538; monogynus, 536–538; opulifolius, 536, 537; pauciflorus, 538; pubescens, 536; Ramaleyi, 536; stellatus, 536, 537
Origin, Some native weeds and their probable, 347
Orobanche, 358
Oscillatoria, 242, 247; Froelichii, 243; Imperator, 243; limosa, 242, 243, 247; princeps, 242, 243, 247; tenuis, 242, 247
Osmunda, 28; cinnamomea, 52; Claytoniana, 52; spectabilis, 52
Oxalis, 352, 358

Oxybaphus nyctagineus, 543
Pachyphyllum araucarinum, 257; rigidum, 257
Pagiophyllum, 257
Pandanus **glauciphyllus,** 64, 74; polycephalus, 65
Pandorina, 241, 246; Morum, 246
Panicularia acutiflora, 197; americana, 196; canadensis, 196; fluitans, 196; nervata, 196; pallida, 196; **septentrionalis,** 196
Peridermium, 25, 506; germinale, 506
Periodicity in certain ponds and streams, Algal, 223
Peronospora, 363, 364; Alsinearum, 543; alta, 552; arborescens, 363, 364, 551; Arenariae macrospora, 543; Arthurii, 551; calotheca, 553; candida, 552; Claytoniae, 552; Corydalis, 363, 364, 550, 551; Cynoglossi, 545; Cyparissiae, 363; Dipsaci, 550; dubia, 364; Echinospermi, 545; effusa, 548; Euphorbiae, 363, 550; Ficariae, 552; Floerkeae, 364, 551; Giliae, 552; grisea, 553; Hedeomae, 551; Hydrophylli, 551; Hyoscyami, 364; leptosperma, 546; Linariae, 553; Lini, 551; Lophanthi, 551; Myosotidis, 545; Nicotianae, 364, 365, 553; Oxybaphi, 543; parasitica, 545, 546; phloginia, 552; Plantaginis, 552; Polygoni, 363, 552; Potentillae, 552, 553; Rumicis, 363; Rubi, 553; Schleideni, 543; Seymourii, 553; sordida, 553; sparsa, 553; trichomata, 544; Trifoliorum, 550; Urticae, 553; Viciae, 550; Violae, 554; Whippleae, 553
Peronosporales, North American, 361, 543
Petrophytum, 538, 540, 541; caespitosum, 541; elatior, 541
Peziza, 529
Phalaris canariensis, 194; arundinacea picta, 194
Phanerophlebia, 27
Pharbitis bracteata, 99
Pharus **parvifolius,** 301; glaber, 302
Phaseolus lunatus, 550
Phegopteris, 28
Phellorina, 294
Philippine Polyporaceae, 391
Philotria, 457; angustifolia, 460–465; canadensis, 458, 459, 464, 465; linearis, 464, 465; minor, 462, 464, 465; Nuttallii, 461, 464, 465; Planchonii, 462, 464, 465
Philotria Raf., Notes on, 457
Phleum pratense, 191
Phlox divaricata, 552
Phormidium incrustatum cataractorum, 248; tenue, 248
Phragmicoma, 161; affixa, 382, 383; aulacophora, 161; bicolor, 159, chinantlana, 160; polycarpa, 162
Phragmidium Libocedri, 509
Phragmites Phragmites, 195
Phyllachora pomigena, 427
Phyllitis, 28
Phymatodes, 28
Physarum contextum, 578
Physocarpa, 535
Physocarpon, 535
Physocarpum, 535
Physocarpus, 535; michiganensis, 536; missouriensis, 536
Phytolacca decandra, 350
Phytophthora Cactorum, 553; infestans, 553; Phaseoli, 550; Thalictri, 363, 552
Pilocereus divaricatus, 565
Pilularia globulifera, 20
Pinus caribaea, 206; insularis, 392, 405, 410; Pumilio, 57; rigida, 55, 57; sylvestris, 57
Piper, 566, 567; peltatum, 556, 567; peltatum hirtellum, 566; umbellatum, 566
Pisonia longirostris, 65
Pistacia Simaruba, 341
Pisum sativum, 550
Placoheppia, 300
Plantago aristata, 552; lanceolata, 552; major, 352, 552; patagonica aristata, 552; Rugelii, 352, 552; virginica, 552
Plants of Nantucket, The ferns and flowering, 49, 181, 471
Plants, Studies of West Indian, 337, 561
Plants, The water-storing tubers of, 271
Plasmopara pygmaea, 552
Platanus, 575
Platycrinus hemisphericus, 572
Platylejeunea, 174, 175; Kroneana, 387, 388
Pleonectria, 528, 530; berolinensis, 530
Pleurocarpus mirabilis, 232, 244
Pleurococcus, 246; vulgaris, 246
Poa, 358; annua, 196; compressa, 196; fasciculata, 197; pratensis, 196; serotina, 196; trivialis, 196, 475
Podisoma gymnosporangium clavipes, 506
Podocarpus Blumei, 63; latifolia, 63
Podozamites acutus, 257; lanceolatus, 257
Poecilopteris crenata, 33
Polyalthia, 68; **clusiflora,** 68
Polygala verticillata, 352

Polygonum, 358, 363; aviculare, 352, 363, 552; dumetorum, 363, 552; erectum, 352; Hydropiper, 352; pennsylvanicum, 352; sagittatum, 352; scandens, 352, 363, 552
Polypodium, 28; vulgare, 22
Polyporaceae, Additional Philippine, 391
Polyporus Adami, 402; asper, 392; badius, 402; brunneopictus, 392; caperatus, 392; carneo-niger, 402; celebicus, 402, 403; coracinus, 403, Dybowski, 399; elegans, 403; fissus, 403; flabelliformis, 403; grammocephalus, 401, 403; Hasskarlii, 412; lucidus, 24; ochrocroceus, 411; palensis, 403; Perula, 402, 403; Philippinensis, 401; rasipes, 402; squamaeformis, 392; strigatus, 392; xeranticus, 393
Polystichum, 28
Polystictus malacensis, 400; malaiensis, 400; Möllerianus, 402; xerampelinus, 397
Pontederia, 58
Populus tremuloides, 532
Porella, 226
Portulaca oleracea, 552
Potamogeton, 224, 236; diversifolius, 59; foliosus, 209, 214–216; lucens, 213–215; mysticus, 59; natans, 209, 214–217; Nuttallii, 59; Oakesianus, 59; pauciflorus, 214; pectinatus, 59; perfoliatus, 59; pulcher, 59; pusillus, 59
Potamogeton lucens, The development of the embryo-sac and embryo of, 209
Potentilla, 358; grandiflora, 553; monspeliensis, 553; nepalensis, 553; norvegica, 553; simplex, 352
Potomorphe, 566, 567; peltata, 566; umbellata, 566
Prenanthes altissima, 549
Prionolejeunea, 374, 375
Protococcus gigas, 235
Protodammara, 249, 254
Prunella vulgaris, 352
Pseudo-araucaria, 254
Pseudoperonospora celtidis, 553; cubensis, 549, 550
Pteridium, 28; aquilinum, 52; aquilinum latiusculum, 52
Pteris, 28
Ptilimnium capillaceum, 352
Ptychanthoides, 161, 162
Ptychanthus, 178
Ptychocoleus, 155, 158, 161, 162, 165, 168; aulacophorus, 161; densifolius, 159; polycarpus, 161, 162, 164, 165, 179; torulosus, 165
Puccinellia, 197; airoides, 475; distans, 197, 475; fasciculata, 475; maritima, 197
Puccinia Botryapites, 506; clavipes, 506
Puerto Rico, Hepaticae of, 155
Putoria, 203
Pycnolejeunea Schwaneckei, 173
Pycnoporus sanguineus, 404
Pyropolyporus albomarginatus, 411; caliginosus, 412; endotheius, 412; fastuosus, 412; lamaensis, 412; pectinatus, 412; **subextensus,** 413; **tenuissimus,** 413; **tricolor,** 413; Williamsii, 412
Pyrus, 505; arbutifolia, 505

Quamoclit Quamoclit, 549; vulgaris, 549

Races, The genus Ernodea Swartz: a study of species and, 203
Radula Grevilleana, 382
RAMALEY, F., & DODDS, G. S. Two imperfectly known species of Crataegus, 577
Ranunculus, 358; abortivus, 352, 552; acris, 452, 552; bulbosus, 452, 552; fascicularis, 552; pennsylvanicus, 552; recurvatus, 552; repens, 552; septentrionalis, 552
Raphanus sativus, 546
Resolutions adopted by the Torrey Botanical Club and other scientific organizations in relation to the death of Lucien Marcus Underwood, 41
Rhus typhina, 352
Rhysotheca australis, 550; Geranii, 551; Gonolobi, 544; Halstedi, 544, 546–548; illinoiensis, 553; obducens, 545; ribicola, 551; Umbelliferarum, 544; Viburni, 546; viticola, 554
Ribes albinervium, 551; divaricatum, 551; glandulosum, 551; hirtellum, 551; oxyacanthoides, 551; prostratum, 551; rotundifolium, 551; rubrum subglandulosum, 551; triste, 551
Riccia, 23, 24
Riella, 30
Rigidoporus surinamensis, 404
Rivea bracteata, 99
ROBINSON, C. B. Alabastra philippinensia, 63
Roestelia, 499, 501, 502, 504–506, 508; aurantiaca, 506; Botryapites, 506; cornuta, 505, 507, 508; Ellisii, 506; flaviformis, 504; floriformis, 504; penicillata, 501, 505; transformans, 505
Roripa, 546; Armoracia, 546; hispida, 546; obtusa, 546; palustris, 546; sessiliflora, 546; sinuata, 546; Walteri, 546

Rosa, 358; californica, 553
Rosaceae, Notes on, 535
Rubus, 352, 354, 356, 358, 360; Baileyanus, 553; canadensis, 553; nigrobaccus, 553; occidentalis, 553; procumbens, 553; strigosus, 553; triloba, 548; villosus, 553; villosus humifusus, 553
Rudbeckia fulgida, 548; laciniata, 548
Rumex, 363; Acetosella, 352
Ruppia maritima, 59, 60
RYDBERG, P. A. Notes on Philotria Raf., 457; Notes on Rosaceae, 535
Rynchospora alba, 483; glomerata, 483; jubata, 569; Torreyana, 483

Sabia **philippinensis** 70, 74; Swinhoei, 71
Sabina, 501; barbadensis, 509; bermudiana, 509; monosperma, 510; pachyphlaea, 510, 511; sabinoides, 508, 511; scopulorum, 510, 511; utahensis, 510, 511; virginiana, 499, 501, 503, 508–511
Sagittaria, 214; Engelmanniana, 62; lancifolia, 213, 217; latifolia, 62, 213; variabilis, 213, 215, 217
Salix, 358; discolor, 352
Salvia lanceolata, 551
Sambucus canadensis, 352
Sarcopyramis **delicata**, 72, 75
Sarothra gentianoides, 352
Sassafras, 571, 573, 575; Sassafras, 574
Sassafras, An analogy between the development of the plates of crinoids and the leaves of, 571
Saururus cernuus, 213, 217
Savastana odorata, 190
Saxifraga, 358
Scenedesmus, 241, 247; caudatus, 247; obtusus, 247
Scientific organizations in relation to the death of Lucien Marcus Underwood, Resolutions adopted by the Torrey Botanical Club and other, 41
Scirpus americanus, 480; cyperinus, 481, 482; cyperinus condensatus, 482; cyperinus pelius, 482; nanus, 480; occidentalis, 481; paludosus, 481; pedicellatus, 482; rubrotinctus, 481; robustus, 481; validus, 481
Schizachyrium scoparium, 181, 182; **littorale**, 182; villosissimum, 182
Schizaea, 29
Schizodendron, 250
Schizostipae, 378
Schoenocrambe linifolium, 546
Scleria triglomerata, 483
Scleroderma strobilinum, 294
Sclerospora Farlowii, 552; graminicola, 552
Scrophularia, 553; californica, 553; marylandica, 553; nodosa, 553; nodosa marylandica, 553
SEATON, S. The development of the embryo-sac of Nymphaea advena, 283
SEAVER, F. J. Color variation in some of the fungi, 307; North Dakota slime-moulds, 577; Some North Dakota Hypocreales, 527
Selaginella, 23, 29, 30; rupestris, 26
Senecio, 548; aureus, 548; aureus croceus, 548; cymbalarioides, 548; integrifolia, 548; Hartianus, 548; oblanceolatus, 548; lugens, 548; Mac Dougalii, 548; peninsularis, 548; Serra, 548; vulgaris, 548
Sequoia gigantea, 293
Serpicula angustifolia, 457; canadensis, 465; occidentalis, 457, 460, 463, 465; verticillata, 457, 459, 465; verticillata angustifolia, 459, 465
Setaria, 552
SETCHELL, W. A. Notes on Lycoperdon sculptum Harkness, 291
Sicyos angulatus, 550
Silene, 358, 543; antirrhina, 352, 543
Silphium integrifolium, 548; laciniatum, 548; perfoliatum, 548; terebinthinaceum, 548; trifoliatum, 548
Sinapis alba, 545, 546; arvensis, 546
Sisymbrium, 546; canescens, 546; incisum, 546; officinale, 546
Sisyrinchium, 353, 358
Slime-moulds, North Dakota, 577
Smilax, 358; rotundifolia, 353
Solanum Lycopersicon, 553; tuberosum, 553
Solidago, 548; canadensis, 352, 548; nemoralis, 352; Riddellii, 548; rigida, 548
Some Araucarian remains from the Atlantic coastal plain, 249
Some Echinocerei of New Mexico, 77
Some native weeds and their probable origin, 347
Some North Dakota Hypocreales, 527
Sonchus, 549; asper, 549
Sophia, 546; Hartwegiana, 546; incisa, 546; pinnata, 546
Sorbus, 508
Sorghastrum avenaceum, 182
Sorosporium, 147; Aristidae, 148; **confusum**, 148; Ellisii, 147, 148; Ellisii Andropogonis, 148; Ellisii Aristidae, 148
Sorosporium Ellisii Winter, a composite species, 147
Sparganium, 215, 217; americanum, 58;

androcladum, 58; eurycarpum, 58; lucidum, 58
Spartina caespitosa, 194; cynosuroides, 194; glabra alterniflora, 194; glabra pilosa, 194; juncea, 194; patens, 194; polystachya, 194
Species, New or noteworthy, 361
Species of Closterium, The New England, 109
Species of Crataegus, Two imperfectly known, 577
Species of Evolvulus, Two Bahamian, 89
Species, Sorosporium Ellisii Winter, a composite, 147
Species and races: The genus Ernodea Swartz, a study of, 203
Specularia perfoliata, 352
Spergula arvensis, 543
Sphaeropsis malorum, 435
Sphenopholis palustris, 475; palustris flexuosa, 475
Spilocaea Pomi, 423
Spinacia oleracea, 548
Spiraea, 538, 540; alba, 539; alleghanensis, 541; arbuscula, 539; Aruncus americana, 541; Beauverdiana, 539; betulifolia, 539; caroliniana, 537; chamaedryfolia, 538; cinerascens, 541; corymbosa, 538; densiflora, 539; Douglasii, 540; Douglasii × densiflora, 540; Douglasii × lucida, 540; Hartwegiana, 538; Helleri, 539, 540; japonica, 539; latifolia, 539, 540; lucida, 538; Menziesii × densiflora, 540; Menziesii × lucida, 540; monogyna, 538; Nobleana, 540; obovata, 540; opulifolia, 535; opulifolia pauciflora, 538; parvifolia, 538; pauciflora, 538; pyramidata, 540; ribifolia, 536; roseata, 540; salicifolia, 539, 540; splendens, 538, 539; Steveni, 538, 539; subcanescens, 540; subvillosa, 540; tomentosa × alba, 540; tomentulosa, 540
Spirodela polyrrhiza, 353
Spirogyra, 226, 228, 230, 231, 233, 244; crassa, 230, 244; inflata, 230, 244; neglecta, 244; nitida, 226–228, 230, 244; rivularis, 244; varians, 227–230, 244; Weberi, 231, 244
Sporobolus vaginaeflorus, 191, 353
Sporogonium of Marchantia polymorpha, The development of the sexual organs and, 321
STANDLEY, P. C. Some Echinocerei of New Mexico, 77
Staurastrum, 109, 234, 245
Stegnosperma, 567; cubense, 567; halimifolium, 567
Stemonitis maxima, 578; Smithii, 578
Stenochlaena, 32, 33
Stichococcus, 237, 238, 246
Stipa avenacea, 191
Strobilomyces, 524
Structure in Carica Papaya, Correlation of flower- and fruit-, 141
Studies in North American Convolvulaceae, 97
Studies in North American Peronosporales, 361, 543
Studies in the genus Gymnosporangium, 499
Studies of West Indian plants, 337, 561
Study of species and races: The genus Ernodea Swartz, 203
Symbiezidium, 388; barbiflorum, 387, 388; **laceratum**, 386–389; transversale, 386; vincentinum, 387, 388
Synchytrium, 423
Syntherisma filiforme, 183, 353; humifusum, 183; sanguinale, 183, 353

Terebinthus, 341; **angustata**, 341, 342; **glauca**, 341, 342; **Hollickii**, 341; **inaguensis**, 341, 342; **Nashii**, 341, 342; Simaruba, 341; **simplicifolia**, 341, 342
Tetraspora, 241, 247; lubrica, 229, 241, 247
Thalictrum **philippinense**, 65, 74; purpurascens, 552
Thelypodium pinnatifidum, 546
Theobroma Cacao, 553
Thesium psilotoides, 65
Thlaspi glaucum, 546; Nuttallii, 546
Thysananthus, 178
Tilia americana, 578
Tilmadoche viridis, 578
Tiniaria dumetorum, 552; scandens, 552
Tissa leucantha, 543; marina, 543
Todea, 29
Torrey Botanical Club and other scientific organizations in relation to the death of Lucien Marcus Underwood, Resolutions adopted by the, 41
Trachylejeunea, 373–375; Aquarius, 374; **dilatata**, 372–375, 389; prionocalyx, 374, 375; Spruceana, 374
Tragopogon porrifolius, 549
Trametes conchata, 405; **conglobata**, 405; **insularis**, 405; Mulleri, 405
Tremella Botryapites, 506; clavipes, 506
Trianthema Portulacastrum, 361, 543
Trichilia **Harrisii**, 568; persimilis, 579
Trichodium laxiflorum, 192
Trichomanes, 29; Petersii, 25
Trichosanthes Anguina, 550, columbrina, 550
Tricuspis sesslerioides, 195
Tridens flava, 195
Trifolium, 358, 550; carolinianum, 550

Triglochin, 215; maritima, 62
Trillium, 25
Triplasis purpurea, 195
Tubers of plants, The water-storing, 271
Tubifera ferruginosa, 579
Two Bahamian species of Evolvulus, 89
Two imperfectly known species of Crataegus, 577
Two new grasses from the West Indies, 301
Tylodendron, 250
Typha, 224, 232, 235, 238; angustifolia, 58; latifolia, 58; orientalis, 63
Tyromyces **Merrittii,** 406; **subchioneus,** 406; **unguliformis,** 406
Udora, 457, 458; canadensis, 459–461, 465; verticillata minor, 462, 463, 465
Ullmannia, 250
Ulothrix, 237, 246; aequalis, 237, 246; subtilis, 246; zonata, 238, 246
Underwood, Lucien Marcus, biographical sketch of, 1; A memorial tribute, 13; Resolutions adopted by the Torrey Botanical Club and other scientific organizations in relation to the death of, 41; The published work of, 17; Relation to the work of the New York Botanical Garden, 39
Unona clusiflora, 68
Uredo nootkatensis, 501
Urtica, 530; gracilis, 352, 553
Urticastrum divaricatum, 553
Uvaria **rubra,** 68, 74; **scandens,** 69, 74
Vallisneria spiralis, 62
Variation in some of the fungi, Color, 307
Vaucheria, 238, 240, 241, 243, 246; geminata, 235, 246; geminata racemosa, 240, 241; sessilis, 240, 246
Verbena urticaefolia, 352
Verbesina encelioides, 548
Vernonia Baldwinii, 548; noveboracensis, 548
Veronica, 553; alpina, 553; Anagallis, 553; Anagallis-aquatica, 553; arvensis, 553; peregrina, 553; Wormskioldii, 553
Vibrio acerosus, 121; Lunula, 122
Viburnum acerifolium, 546; dentatum, 546; nudum, 546; Opulus, 546; pubescens, 546
Vicia, 550; americana, 550; americana linearis, 550; Faba, 550; linearis, 550; sativa, 550
Vincetoxicum, 544; gonocarpos, 544; hirsutum, 544; suberosum, 544
Viola, 352, 354, 358, 486; odorata, 554; Rafinesquii, 554; tenella, 554; tricolor, 554; tricolor arvensis, 554
Vitis aestivalis, 554; bicolor, 554; californica, 554; cinerea, 554; cordifolia, 554; Labrusca, 554; riparia, 554; rotundifolia, 554; vinifera, 554; vulpina, 554
Vittaria, 29
Voltzia, 250

Walchia, 250
Water-storing tubers of plants, 271
Wedelia incarnata, 543
Weeds and their probable origin, Some native, 347
West Indian Lejeuneae, New, 371
West Indian plants, Studies of, 337, 561
West Indies, Two new grasses from the, 301
WESTER, P. J. The correlation of flower- and fruit-structure in Carica Papaya, 141
Whipplea modesta, 553
Whitfordia, 407; **Warburgiana,** 407
WILSON, G. W. Studies in North American Peronosporales, 361, 543
Woodsia, 29
Woodwardia, 29; paradoxa, 33
Work of Lucien Marcus Underwood, The published, 17

Xanthium, 352; canadense, 544
Xanthoria, 299; **modesta,** 298

ZAHLBRUCKNER, A. New North American lichens, 297
Zannichellia, 215–217; palustris, 60, 214
Zanthoxylum americanum, 532
Zizania aquatica, 190
Zostera marina, 62
Zygnema, 231, 244; cruciatum, 231, 244; leiospermum, 231, 244; insigne, 231, 244